

University of Groningen

## Isomerisatie van nitrophenylsulfonylureumderivaten

Groot, Jacob

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1949

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Groot, J. (1949). *Isomerisatie van nitrophenylsulfonylureumderivaten*. Noordhoff Uitgevers.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.


### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## SUMMARY.

This thesis describes the preparation of derivatives of nitrophenylsulphonyl urea and isourea and the decomposition of the o- and p-nitro compounds under the influence of alkali. This decomposition is an intramolecular isomerization followed by loss of sulphur dioxide:



The derivatives of isourea, when heated with alkali, also lose sulphur dioxide, but moreover the ether function is disrupted and urea derivatives are formed:



This hydrolysis is also known for O,N,N-trimethylisourea, which compound is decomposed by boiling water into dimethylurea and methyl alcohol.

It is probable that the derivatives of nitrophenylsulphonylisourea first isomerise with loss of sulphur dioxide, after which the isourea derivative formed decomposes into a derivative of urea and methyl alcohol. This supposition is confirmed by the fact that the isomerization of N-p-nitrophenylsulphonyl-O-methylisourea is faster than that of p-nitrophenylsulphonylurea. This would not be the case, should the first step be the formation of a sulphonylurea derivative.

On p. 50 a survey is given of all the compounds for which the behaviour in the presence of hot alkali has been examined; the second column mentions the decomposition products.

The compounds with a meta nitro group do not undergo an isomerization. The presence of an o- or p-nitro group is necessary to weaken the bond between the sulphonyl group and the benzene nucleus.

In the presence of an excess of alkali the isomerization is a monomolecular reaction. A reaction mechanism is proposed (p. 53); the accelerating influence of the methyl and phenyl groups is ascribed to a reduction of the activation energy, necessary for the rupture of the carbon sulphur bond. Indeed it was found that this energy is diminished by the introduction of these groups.